| Agent | Degree of evidence of carcinogenicity ^{<i>a,b</i>} | | Overall evaluation of carcinogenicity - to humans ^a |
|---|---|---------------|--|
| | Humans | Animals | |
| Aldicarb | ND | I | 3 |
| Atrazine | I | L | 2B ^c |
| Captafol | ND | S | 2A ^c |
| Chlordane | Ι | S | 2B |
| DDT | I | S | 2B |
| Deltamethrin | ND | I | 3 |
| Dichlorvos | I | S | 2B |
| Fenvalerate | ND | I | 3 |
| Heptachlor | I | S | 2B |
| Monuron | ND | L | 3 |
| Occupational exposures in spraying and application of nonarsenical insecticides | L | | 2A |
| Pentachlorophenol | Ι | S | 2B |
| Permethrin | ND | Ι | 3 |
| Picloram | ND | L (techgrade) | 3 |
| Simazine | I | I | 3 |
| Thiram | I | I | 3 |
| Trifluralin | I | L (techgrade) | 3 |
| Ziram | ND | L | 3 |

SUMMARY OF FINAL EVALUATIONS

^{*a*}I, inadequate evidence; S, sufficient evidence; L, limited evidence; ND, no data ^{*b*}For definitions of degrees of evidence and groupings of evaluations, see Preamble, pp. 26-28. ^cOther relevant data influenced the making of the overall evaluation.