# SUPPLEMENTARY CORRIGENDA TO VOLUMES 1–77

## Volume 63

#### **Benzofuran monograph**

p. 436:

2nd-3rd lines of last paragraph: delete and chromosomal aberrations.

p. 437:

Last line of table should read:				
CIC, Chromosomal aberrations, Chinese	_	-	280	US National Toxicology
hamster ovary (CHO) cells in vitro				Program (1989)

### p. 438

Last paragraph of section 5.4 Other relevant data should read:

Induction of gene mutation and sister chromatid exchange was seen in cultured rodent cells treated with benzofuran in single studies. Benzofuran was not mutagenic to bacteria.

#### Volume 69

n	50'	7.
P٠	59	<i>'</i> .

Reference Rappe et al. (1978b) should read:

Rappe, C., Garå, A. & Buser, H.-R. (1978b) Identification of polychlorinated dibenzofurans (PCDFs) in commercial chlorophenol formulations. *Chemo-sphere*, 7, 981–991

Reference Rappe et al. (1979b) should read:

Rappe, C., Buser, H.-R., Kuroki, H. & Masuda, Y. (1979b) Identification of polychlorinated dibenzofurans (PCDFs) retained in patients with Yusho. *Chemosphere*, 4, 259–266

#### Volume 71, Part 1

p. 233, first line of 2nd paragraph of Section 2.2 Cohort studies, *replace* Shouqui *et al.* (1989) *by* Li *et al.* (1989).

p. 249:

Reference Shouqi et al. (1989) should read:

Li, S.Q., Dong, Q.N., Liu, Y.Q. & Liu, Y.G. (1989) Epidemiologic study of cancer mortality among chloroprene workers. *Biomed. Environ. Sci.*, 2, 141–149

#### IARC MONOGRAPHS VOLUME 78

## Volume 74

p. 111, line 7, sentence should read: 'Thus, depleted uranium has a lower specific activity (12 200 Bq/g) than natural uranium (25 900 Bq/g), and its chemical toxicity predominates over the radiation effects when it is taken into the body in a soluble form (Agency for Toxic Substances and Disease Registry, 1998).'

562